

Document downloaded from:

<http://hdl.handle.net/10251/73485>

This paper must be cited as:

Penella, C.; González Nebauer, S.; Quinones, A.; San Bautista Primo, A.; López Galarza, SV.; Calatayud, A. (2015). Some rootstocks improve pepper tolerance to mild salinity through ionic regulation. *Plant Science*. 230:12-22. doi:10.1016/j.plantsci.2014.10.007.



The final publication is available at

<https://dx.doi.org/10.1016/j.plantsci.2014.10.007>

Copyright Elsevier

Additional Information

1  
2 **Some rootstocks improve pepper tolerance to mild salinity through ionic**  
3  
4 **regulation**  
5  
6  
7

8  
9 **Corresponding author:**  
10

11 Dr. Angeles Calatayud  
12

13 Instituto Valenciano de Investigaciones Agrarias (IVIA). Departamento de  
14 Horticultural. Ctra. Moncada-Naquera km. 4.5. 46113-Moncada, Valencia,  
15  
16  
17  
18  
19 Spain.  
20

21  
22 Phone: +34 963424039  
23

24  
25 Fax. +34 963424000  
26

27  
28 e-mail: calatayud\_ang@gva.es  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 **Some rootstocks improve pepper tolerance to mild salinity through ionic**  
3 **regulation**  
4  
5

6  
7 Consuelo Penella<sup>a</sup>, Sergio G. Nebauer<sup>b</sup>, Ana Quiñones<sup>a</sup>, Alberto San Bautista<sup>b</sup>,  
8  
9 Salvador López-Galarza<sup>b</sup>, Angeles Calatayud<sup>a,\*</sup>  
10  
11

12  
13  
14  
15 <sup>a</sup>Instituto Valenciano de Investigaciones Agrarias (IVIA). Departamento de  
16  
17 Horticultural. Ctra. Moncada-Naquera km. 4.5. 46113-Moncada, Valencia,  
18  
19 Spain.  
20  
21

22  
23 <sup>b</sup>Universitat Politècnica de València. Departamento de Producción Vegetal.  
24  
25 Camino de Vera 14, 46020 Valencia, Spain.  
26  
27

28  
29  
30  
31 \*Corresponding author: Angeles Calatayud<sup>a</sup>. Instituto Valenciano de  
32  
33 Investigaciones Agrarias (IVIA). Departamento de Horticultural. Ctra. Moncada-  
34  
35 Naquera km. 4.5. 46113-Moncada, Valencia, Spain. e-mail:  
36  
37 calatayud\_ang@gva.es  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 **ABSTRACT**  
3

4 Grafting has been proposed as an interesting strategy that improves the  
5 responses of pepper cultivars under salinity. However, very little is known  
6 about the physiological mechanisms underlying the increased tolerance  
7 provided by rootstocks on the scions. With this aim, we performed this  
8 experiment. The commercial 'Adige' pepper cultivar was grafted onto three  
9 different pepper accessions that showed differential salt tolerance (accessions  
10 5, 12 and 14). Responses to salinity (40mM NaCl) were studied for 30 days by  
11 determining water relations, mineral content, proline accumulation,  
12 photosynthetic parameters, nitrate reductase activity and antioxidant  
13 capacity. The responses observed were depended on salinity treatment  
14 duration and the rootstock used. Higher salt tolerance was achieved when the  
15 'Adige' cultivar was grafted onto the 12 genotype, which allowed not only lower  
16 Na<sup>+</sup> and Cl<sup>-</sup> accumulation in the scion, but also ion selectivity maintenance,  
17 particularly Na<sup>+</sup>/K<sup>+</sup> discrimination. These traits led to a minor negative impact on  
18 photosynthesis, nitrate reductase activity and lipid peroxidation in grafted scion  
19 leaves. This work suggests that using tolerant pepper rootstocks that maintain  
20 the scion's ion homeostasis is a promising strategy to provide salinity tolerance  
21 and can consequently improve crop performance when faced with farmland  
22 salinity.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 *Key words:* Graft; NaCl; Ions; Pepper; Photosynthesis; Water relations  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

Grafting plants onto tolerant rootstocks is one of several approaches that can reduce the impact of salinity [1], one of the most serious problems of horticultural crops in arid and semi-arid regions [2].

Pepper is one of the most important vegetable crops in these areas and is considered sensitive to salinity [3], even though salt tolerance can vary between pepper genotypes [4]. Some pepper accessions have been identified as salinity-tolerant and have been successfully used as pepper rootstocks under saline conditions [5].

Several studies have been conducted in tomato and melon to elucidate the mechanisms involved in increased salinity tolerance of grafted plants. This increased tolerance of grafted plants is generally associated with their capacity to exclude or retain and/or accumulate toxic ions,  $\text{Na}^+$  and  $\text{Cl}^-$  in rootstock roots, thus limiting their transport to leaves rather than through the synthesis of osmotically active metabolites or the induction of antioxidant systems [6–8]. Other authors have indicated that influence of rootstock on the salt tolerance of the scion is due to a more efficient control of stomatal functions (changes in stomatal regulation and water relations), which indicate that the grafting incision may alter hormonal signalling between roots and shoots [9]. In other cases, this raised tolerance has been explained by the re-establishment of ionic homeostasis [10].

Nevertheless, the mechanism of resistance against salinity in grafted plants displays great complexity in association with specific rootstock/scion interactions [11,12], and can vary among species. As far as we know, very few studies of this type have been conducted in pepper to elucidate whether or not

1 salt tolerance conferred by rootstocks is also due to exclusion and/or retention  
2 mechanisms, as in tomato or melon given their better capacity to alleviate the  
3 toxic effects of salts or other processes; e.g., maintenance of water relations or  
4 antioxidant capacity. Guifrida et al. [13] found that stunted growth due to salinity  
5 was attenuated in pepper-grafted plants when compared to non-grafted plants  
6 associated primarily with reduced uptake of salt ions and, therefore, with a  
7 lower concentration of these ions in the grafted plants instead of maintaining  
8 leaf turgor by osmotic adjustments.  
9

10 To answer this question, in previous experiments we selected three  
11 pepper accessions with different degrees of salinity tolerance [5] under mild salt  
12 stress. In this study, we aimed to identify the physiological responses to salinity  
13 stress involved in increased tolerance of pepper-grafted plants using these  
14 accessions as rootstocks and to elucidate if mechanisms to tolerance are  
15 related with the role of roots rootstocks in altering the stress perception by the  
16 scion. To fulfil these objectives, we discussed differences in pepper-grafted  
17 plants adaptation mechanisms in response to mild salt stress by comparing  
18 some physiological parameters: photosynthesis; lipid peroxidation levels;  
19 relative water content (RWC); proline concentration; osmotic potential ( $\Psi_s$ ); ions  
20 concentration; nitrate reductase activity (NR). We present evidence that grafting  
21 plants onto appropriate (tolerant) rootstocks is a good tool against salinity  
22 stress, which is mediated mainly by reducing ionic toxicity to the scion.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 **2. Materials and methods**

### 55 *2.1. Plant material and greenhouse conditions*

56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Based on previous studies, we selected three pepper accessions (wild types) with a different salinity tolerance [5]: 'ECU-973' of *Capsicum chinense* Jacq. (code 12) as being tolerant; 'BOL-58' of *Capsicum baccatum* L. var. *pendulum* (code 14) as being moderately tolerant; and 'Serrano' of *Capsicum annuum* L. (code 5) as being less tolerant. These accessions were chosen as rootstocks and the pepper cultivar 'Adige' (Lamuyo type, Sakata Seeds, Japan) was grafted onto these three pepper accessions in this study. Pepper seeds were sown on 1 December in 100-cell polystyrene trays filled with peat-based substrate and kept in a Venlo-type glasshouse. The graft was performed on 12 February using the tube-grafting method (cutting the growing tip of the rootstock at a 45° angle below the cotyledons, attaching to the scion, previously cut at a 45° angle above the cotyledons, and fixing the rootstock and scion with a clip).

One month after grafting, the root system of the plants was washed to clean the substrate and plants were placed in 5 L polyethylene pots covered with aluminium sheets. Pots were filled with a standard nutrient solution for pepper [14]. The electrical conductivity (EC) and pH of this nutrient solution was 1.7 dS m<sup>-1</sup> and 6.5, respectively. Nutrient solution was added daily to compensate for uptake. After 7 days of leaving seedling plants to acclimatise to pots, salinity treatment was initiated by adding NaCl (40mM) to the nutrient solution to reach an EC of 5.2 dS m<sup>-1</sup> NaCl.

Treatments were defined by two salinity levels (0 and 40mM NaCl) and four plant combinations: the cultivar 'Adige' grafted onto rootstock accessions 5, 12 and 14, and ungrafted 'Adige' plants were used as the controls.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The grafted combinations (cultivar/rootstock) were labelled as A/5, A/12 and A/14. The layout was completely randomised with three replications per combination and six plants per replication.

All the physiological measurements were taken on 14 (T1) and 28 (T2) days after NaCl addition on fully expanded mature leaves (third or fourth leaf from the shoot apex).

During the culture, plants were grown in a Venlo-type greenhouse under natural light conditions ( $610\text{-}870\ \mu\text{mol m}^{-2}\ \text{s}^{-1}$ ), temperature ranges were  $21\text{-}24^{\circ}\text{C}$ , and relative humidity was  $52\text{-}72\%$ .

## 2.2. *Water relations*

The osmotic potential of leaf sap ( $\Psi_s$  in MPa) was measured with an osmometer (Digital osmometer, Wescor, Logan, USA). Two independent determinations were made on each replicate and plant combination, obtained from six plants per treatment and combination at T1 and T2.

Leaves were tightly wrapped in aluminium foil, frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$ . After thawing, sap was collected from syringes at  $25^{\circ}\text{C}$  and placed in the osmometer. Osmolyte content ( $\text{mmol kg}^{-1}$ ) was converted into MPa using the Van't Hoff equation [15].

Six other similar leaves from two independent plants of each plant combination, salinity treatment and replicate were collected to determine the (RWC) as  $(\text{FW}-\text{DW})/(\text{TW}-\text{DW}) \times 100$ , where FW is fresh weight, DW is dry weight, and TW is turgid weight [15].



### 2.3. Ion analysis

The leaves and roots collected at T1 and T2 for  $n \geq 5$  samples of each treatment and plant combination were dried at 70°C for 4 days. Dried samples were digested in a mixture at 70% of HNO<sub>3</sub>-HClO<sub>3</sub> (2:1). Macronutrients (K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup> and Na<sup>+</sup>) were measured by ICP emission spectrometry (iCAP 6000, Thermo Scientific. Cambridge, United Kingdom).

The chloride concentration (Cl<sup>-</sup>) in the dry plant material was extracted with 0.1N HNO<sub>3</sub> in 10% (v/v) acetic acid and was determined by potentiometric titration with AgNO<sub>3</sub> in a chloride analyzer (Sherwood, MKII 926). The results were expressed as mg g<sup>-1</sup> DW.

### 2.4. Proline determination

Proline content (mg g<sup>-1</sup> DW) was determined as described by [16]. Leaf pepper tissue (0.05 g) was ground in 3% sulphosalicylic acid, the homogenate was filtered, and 0.75 mL of glacial acetic acid and 0.75 mL of ninhydrin reagent (1.25 g ninhydrin in 30 mL glacial acetic acid and 20 mL 6N phosphoric acid) were added to an aliquot of the filtrate. The reaction mixture was boiled for 1 h, and readings were taken at a wavelength of 520 nm in a spectrophotometer. Three independent determinations were made in three different extracts obtained from 18 plants per treatment and combination (one leaf per plant, and six plants per extract).

### 2.5. Photosynthetic activity and chlorophyll fluorescence

The CO<sub>2</sub> fixation rate ( $A_N$ ,  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ), stomatal conductance to water vapour ( $g_s$ ,  $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ ), transpiration rate ( $E$ ,  $\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$ ) and

1 substomatal CO<sub>2</sub> concentration (C<sub>i</sub>, μmol CO<sub>2</sub> mol<sup>-1</sup> air) were measured in the  
2 steady state while maintaining plants at 1,000 μmol m<sup>-2</sup> s<sup>-1</sup> for 10-15 min and  
3  
4 400 ppm CO<sub>2</sub> with a LI-6400 (LI-COR, Nebraska, USA). Light curves were  
5  
6 previously performed (data not shown) and A<sub>N</sub> was saturated at 900 μmol m<sup>-2</sup> s<sup>-1</sup>  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1. The gas exchange and fluorescence determinations were made from 9 am to 11 am (GMT). One measurement per plant was taken, and ten different plants were used (n=10) for each treatment (control and salinity stress) and plant combination.

## 2.6. Nitrate reductase activity

Nitrate reductase activity (EC 1.6.6.1) in leaves was determined *in vivo* following the methods described by [17,18]. Discs, 1 cm in diameter, were punched out of mature fresh leaves. Samples (200 mg) were suspended in a glass vial containing 10 mL of 100 mM potassium phosphate buffer (pH 7.5), 1% (v/v) *n*-propanol and 100 mM KNO<sub>3</sub>. The glass vial was subjected 3 times to vacuum infiltration in order to induce anaerobic conditions in the incubation medium. Plant samples were incubated in a water bath at 30°C for 60 min in the dark and were placed in a boiling water bath for 5 min to stop the enzymatic reaction. The nitrite released from the plant material was determined colorimetrically at 540 nm (spectrophotometer PerkinElmer, Lambda 25) by adding 0.02% (w/v) N-naphthylethylenediamine and 1% sulphanilamide. A standard curve with KNO<sub>2</sub> was prepared to calculate the amount of NO<sub>2</sub> that the samples contained. Sampling and replicates were used as described for proline determination.

## 2.7. Lipid peroxidation

Lipid peroxidation in leaves was estimated through malondialdehyde (MDA) determinations using the thiobarbituric acid reaction following the protocol reported by [19], and modified in [20]. The non-specific background absorbance reading at 600 nm was subtracted from the specific absorbance reading at 532 nm. The sampling and replicates used were those described for proline determination.

## 2.8. Statistical analyses

The results were subjected to a variance analysis (ANOVA; Statgraphics Centurion for Windows, Statistical Graphics Corp.). The mean comparisons were made using Fisher's least significance difference (LSD) test at  $P < 0.05$ . The data obtained in some measurement parameters were subjected to linear regression and analyses to identify the relationships between the parameters.

# 3. Results

## 3.1. Water relations

Plant water relations were assessed by the determination of RWC and  $\Psi_s$  (Figs. 1 and 2). No changes in RWC were observed in the experiment in any plant combination, except for ungrafted plants (Fig. 1A, B), where RWC diminished ( $P < 0.05$ ) after salt treatment.

The  $\Psi_s$  of all the plant combinations reduced significantly ( $P < 0.05$ ) under salinity at T1 and T2 (Fig. 2). At T1, no significant interaction was found. At T2, differences between treatments were greater in ungrafted and A/5 than in A/12 and A/14 ( $P < 0.05$ ).

### 3.2. Ion partitioning

The Na<sup>+</sup> concentration in leaves and roots increased under NaCl (Fig. 3A) in all the plant combinations. The Na<sup>+</sup> concentration in leaves was higher in ungrafted and A/5 plants (Fig. 3A) if compared with A/12 and A/14 (P<0.05) at T1 and T2 under salinity. In general terms, the Na<sup>+</sup> concentration in the roots under salinity was higher than in leaves (Fig. 3B), with a lower concentration found in A/12 and A/14.

Chloride content was approximately 4 times higher than Na<sup>+</sup> in leaves. The Cl<sup>-</sup> concentration in leaves (Fig. 3C) increased with a higher NaCl concentration and time exposure, but this incident did not occur in roots (Fig. 3D) and in none of the plant combinations. Ungrafted and A/5 obtained the highest Cl<sup>-</sup> levels in leaves, whereas A/12 and A/14 plants showed a greater accumulation in roots (P<0.05) (Fig. 3D).

In general terms, a consistent K<sup>+</sup> content reduction trend was observed in leaves at T1 under saline conditions in all the plant combinations (Fig. 3E). This decrease occurred at T2 only in ungrafted and A/5 plants, but not in A/12 and A/14, where no significant differences in the K<sup>+</sup> levels were found if compared with their controls (Fig. 3E). In roots, a marked increase in K<sup>+</sup> content was observed in A/12 at T1 (Fig 3F). In contrast, the K<sup>+</sup> concentration at T2 did not change in A/5 and A/14 under salinity (Fig. 3F).

The Na<sup>+</sup>/K<sup>+</sup> ratio increased significantly depending on salt application and the exposure time in the ungrafted and A/5 leaves (Fig. 3G). The lower values (P<0.05) in leaves were observed for 12/cultivar and 14/cultivar. In the root compartment (Fig. 3H) under salt treatment at T1, the Na<sup>+</sup>/K<sup>+</sup> values increased

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

in ungrafted and A/5. At T2, the  $\text{Na}^+/\text{K}^+$  ratio in roots lowered under salt conditions if compared to the values obtained at T1 in these plant combinations due to a sharp drop in the  $\text{Na}^+$  content in roots at T2.

The  $\text{Ca}^{2+}$  (Fig. 4A) and  $\text{Mg}^{2+}$  levels (Fig. 4C) were similar in leaves for the tandem ungrafted and A/5 plants, with reduced plant exposure to NaCl (Fig. 4). In A/12 and A/14, the  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations in leaves showed minor variations between the control and treated samples (Fig. 4 A, C). In roots, the  $\text{Mg}^{2+}$  levels (Fig. 4D) lowered in all the plant combinations with time, while the  $\text{Ca}^{2+}$  levels lowered in A/5 at T1 and T2, but increased in ungrafted, A/12 and A/14 at T2 (Fig. 4B).

### 3.3. *Proline content in leaves*

Under the control conditions, no significant differences were found in the proline leaf content between plant combinations with time. Salinity gave rise to increased leaf proline content ( $P < 0.05$ ). This increase was similar for all the plants at T1 (Fig. 5A). At T2 (Fig. 5B) under 40mM NaCl, proline content substantially increased in ungrafted and A/5 if compared with their control values, but not in 12/cultivar and 14/cultivar ( $P < 0.05$ ), which showed similar values to T1.

### 3.4. *Gas exchange parameters*

As shown in Figure 6, the  $A_N$  (Fig. 6A, B) and  $g_s$  (Fig. 6C, D) of the grafted plants did not differ from those of the ungrafted plants under the control conditions. The photosynthesis rate significantly lowered in all the plants

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

( $P < 0.05$ ) in response to salt stress, except 12/cultivar at T2, when the  $A_N$  values did not significantly differ from those of the control (Fig. 6B).

A decrease in  $g_s$  under salt treatment was observed in all the plants (Fig. 6C, D). Significant differences were found for the ungrafted, A/5 and A/14 plants if compared to 12/A at T1 and T2. A minor decrease, but with a significant difference compared to its control, was noted for 12/cultivar.

Instantaneous carboxylation efficiency, estimated by the  $A_N/C_i$  ratio (Fig. 6E, F), reduced in ungrafted, A/5 and A/14 at T1 and T2. Interestingly at T2, minor differences were seen in the  $A_N/C_i$  values in A/12, followed by A/14, if compared to their controls, but no significant differences were observed between them.

### 3.5. Nitrate reductase activity in leaves

Salt stress resulted in diminished NR activity in leaves after 14 (Fig. 7A) and 28 (Fig 7B) days of mild NaCl treatment. Under salinity, the greatest NR activity at T1 and T2 was seen for A/12 plants, with significant differences ( $P < 0.05$ ) if compared to ungrafted and A/5. Nevertheless, the inhibition percentages due to salt application at T2 were not associated with the NR control values: 74% for ungrafted, 50% for 5/cultivar, 22% for 12/cultivar and 32% for 14/cultivar (Fig. 7B).

### 3.6. Lipid peroxidation

At T1 (Fig. 8A), MDA content increased and significant differences were observed only in the ungrafted plants. After 28 days of salt exposure, lipid peroxidation increased significantly in the ungrafted and 5/cultivar plants

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

( $P < 0.05$ ). It is noteworthy that no further MDA accumulation occurred in any of the plant combinations (Fig. 8B).

### 3.7. *Relationship between osmotic potential, ions and proline concentrations and photosynthesis in leaves*

Regression analyses were performed with the physiological study parameters. Only the significant linear relations that contribute to understanding tolerance mechanisms to salinity ( $\text{Na}^+$ ,  $\text{Cl}^-$  and  $\text{K}^+$  concentration and proline level vs. osmotic potential and  $A_N$ ) are shown in Table 1.

At T1 and T2, the data gave an inverse linear relationship among  $\Psi_s$  and  $\text{Na}^+$ ,  $\text{Cl}^-$  and proline, but a positive correlation with the  $\text{K}^+$  level in leaves. Proline was the parameter that obtained the steepest slope values to modify  $\Psi_s$ .

$A_N$  at T1 correlated negatively with the  $\text{Na}^+$ ,  $\text{Cl}^-$  and proline concentrations, but not significantly only for the last parameter ( $P > 0.05$ ). The regression analysis indicated inhibition of  $A_N$  with greater dependency of  $\text{Na}^+$  and  $\text{Cl}^-$ . Nevertheless at T2,  $A_N$  lowered, which was due mainly to an increased proline concentration. Although  $A_N$  showed a positive dependency with the  $\text{K}^+$  levels, no significant influence was found, not even at T1 and T2.

## 4. Discussion

NaCl addition is associated with differential responses in physiological parameters in ungrafted and grafted pepper plants. We demonstrate that tolerance to moderate salt stress can be improved by grafting. The best salt acclimation was obtained when accession 12 was used as a rootstock (A/12),

1 based on the minor negative effects caused by salt treatment on  
2 photosynthesis, NR activity and lipid peroxidation. Furthermore, some  
3 favourable physiological characteristics for salt acclimation, such as higher K<sup>+</sup>  
4 Ca<sup>2+</sup> and Mg<sup>2+</sup> levels in leaves and a lower Na<sup>+</sup>/K<sup>+</sup> ratio, were seen in this plant  
5 combination. The latter parameter has been demonstrated as a good indicator  
6 of salt tolerance [21].  
7  
8  
9  
10  
11  
12  
13

14 Salt tolerance in plants is usually associated with the ability to restrict the  
15 uptake and/or transport of saline ions from roots to leaves and their  
16 compartmentalisation [22]. In this study, more Cl<sup>-</sup> was withheld in the roots of  
17 rootstocks in A/12 and A/14, and less Cl<sup>-</sup> was transported to their leaves if  
18 compared with the ungrafted and A/5 plants under NaCl stress (P<0.05). This  
19 suggests either maximised Cl<sup>-</sup> retrieval to the rootstock or a retention  
20 mechanism in the roots of these plant combinations. Unlike Cl<sup>-</sup>, rootstocks 12  
21 and 14 showed a reduced Na<sup>+</sup> net uptake, consequently their leaves gave a  
22 lower Na<sup>+</sup> concentration value if compared with the others (P<0.05). Two  
23 mechanisms can explain the lower Na<sup>+</sup> concentration in roots: firstly, as  
24 suggested by Aktas et al. [4], in salt-tolerant pepper genotypes, a plasma  
25 membrane Na<sup>+</sup>/H<sup>+</sup> antiporter protein is activated in root cells upon NaCl  
26 exposure. This mechanism has been reported in different grafted plants, such  
27 as melon [23–25], tomato [22,26,27], watermelon [28] and cucumber [29].  
28 Alternatively, the root system of rootstocks 12 and 14 might be able to control  
29 Na<sup>+</sup> influx, as reported for pumpkin roots [8].  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Regarding concentration; leaf Cl<sup>-</sup> accumulation exceeded that of Na<sup>+</sup> in  
54 all the plant combinations. This is in accordance with the results obtained by  
55 Navarro et al. [30] and Chartzoulakis and Klapaki [31] in the 'Orlando' variety  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 and the 'Sonar' pepper variety, respectively. The higher  $\text{Cl}^-$  concentration, if  
2 compared to  $\text{Na}^+$  (mainly in roots), can be linked to a higher passive uptake root  
3 component and a very feebly active  $\text{Cl}^-$  uptake system [32]. However, it is  
4 unknown whether some rootstocks are capable of regulating the transport of  
5  $\text{Na}^+$  or/and  $\text{Cl}^-$  to leaves [33]. Based on our results, the capacity to regulate  $\text{Na}^+$   
6 and  $\text{Cl}^-$  uptake and transport was linked to the ability of rootstocks, and not to  
7 the grafting process itself (comparing A/5 vs. A/12 and A/14), indicating that the  
8 physiological and biochemical mechanisms of these salts operate at the  
9 rootstock level, as observed in grafted melon plants [7] or cucumber plants [8].

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21 Regulation of ion homeostasis and selectivity, particularly  $\text{Na}^+/\text{K}^+$   
22 discrimination, is closely linked to the lower  $\text{Na}^+$  concentration and its relation to  
23 salt tolerance [34]. Given the similar physico-chemical structure between  $\text{Na}^+$   
24 and  $\text{K}^+$ , a high  $\text{Na}^+$  concentration in the external solution can lower the  $\text{K}^+$  level  
25 in the tissues of many plants species [35]. In our study, the  $\text{Na}^+/\text{K}^+$  ratio in  
26 leaves of the ungrafted and A/5 pepper plants under salinity was significantly  
27 higher ( $P < 0.05$ ) than those of the plants grafted onto rootstocks 12 and 14, and  
28 the latter is able to select, use and transport  $\text{K}^+$  to leaves, as in many vegetable-  
29 grafted plants exhibiting salinity tolerance; e.g., tomato [6], melon [9] or  
30 cucumber [11,29]. However, the direct relation between  $\text{K}^+$  homeostasis and  
31 salinity tolerance has not been well-established [1]. In some species,  $\text{Na}^+$  can  
32 be balanced by a higher  $\text{K}^+$  concentration [36], while in other plants, tolerance is  
33 due to the capacity of roots to maintain  $\text{K}^+$  transport in the xylem, as in tomato-  
34 grafted plants [37,38].

35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Despite the negative effect on plant growth derived from its toxic effect,  
accumulation of ions under salinity can help maintain the turgor pressure of

1 plants [30,39]. In addition, different osmolytes can be involved in the reduction  
2 of  $\psi_s$ , including organic compounds such as sugars, free amino acids,  
3 glycinebetaine, soluble proteins, proline and organic acids [40–42], and/or  
4 macronutrients such as inorganic components [43]. According to our results, a  
5 strong negative correlation between the reduction in leaf  $\psi_s$  and salt ions  
6 content for all the plant combinations was observed in the experiment. The  
7 linear regressions equations showed that  $\text{Na}^+$  and  $\text{Cl}^-$  display a different  
8 response on  $\psi_s$ . The lower osmotic potential seems to be achieved mainly by  
9  $\text{Na}^+$  and, to a lesser extent by  $\text{Cl}^-$ . This can be explained by a more marked  
10 change in  $\text{Na}^+$  accumulation if compared to  $\text{Cl}^-$  between the ungrafted and A/5  
11 vs. A/12 and A/14 plants, rather than by the absolute concentration of both ions.  
12 The reduced osmotic potential assigned to  $\text{Na}^+$  was consistent with pepper  
13 plants [44], and salt-tolerant species such as *Centaurea ragusina* [45], *Atriplex*  
14 *nummularia* [46] or *Aster tripolium* [47]. The contribution of  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  to  
15  $\psi_s$  under the salinity conditions in our study was more relevant in the A/12 and  
16 A/14 plants at T2, where  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  represented 30-35% of the total  
17 ions if compared to 15% in the ungrafted and A/5 plants.

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42 The adjustment of the osmotic potential through inorganic ion uptake  
43 supposes a much lower energy cost than that conferred by the organic  
44 molecules synthesised in the cell [48]. However in order to reduce  $\psi_s$ , our plants  
45 required proline synthesis to produce sufficient osmotics under salt stress  
46 conditions. The synthesis and accumulation of proline depended on plant  
47 combinations and time exposure. At T1, when the ionic-osmotic phase was  
48 predominant, proline accumulation contributed less. In contrast at T2, strong  
49 proline synthesis took place in the ungrafted and A/5 plants when compared  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 with A/12 and A/14 ( $P < 0.05$ ), and as result, the reduction in  $\psi_s$  strongly related  
2 with proline accumulation in these plant combinations ( $r^2 = 0.95$ ), but more  
3 weakly in the latter ones ( $r^2 = 0.36$ ). An larger amount of proline or other  
4 compatible solutes may protect plants by scavenging the oxygen-free radicals  
5 caused by salt stress [1,29], which has been observed in different grafted plants  
6 like tomato [49], cucumber [29] or tobacco [50]. Conversely, the concomitant  
7 increase in proline with a prolonged exposure time in the ungrafted and A/5  
8 plants was consistent with the higher leaf proline concentrations in salt-sensitive  
9 genotypes reported for other species such as wheat [51], barley [52], *Centaurea*  
10 *ragusina* [45] or rice [53]. This indicates that significant proline accumulation  
11 generally occurs only beyond the salt stress threshold [54]. The energy cost  
12 imposed by ion exclusion and/or compartmentalisation mechanisms are  
13 relatively low when compared with the synthesis of organic molecules [52,55]  
14 but, conversely, accumulation of saline ions may interfere with the normal  
15 biochemical activities taking place within the cell [56].

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
Plants respond to lower water availability under salinity by reducing their  
leaf transpiration, stomatal conductance, and by adjusting their root water  
uptake [57]. Under prolonged periods of exposure to salt, root conductivity can  
be partially recovered, mainly through the accumulation of compatible solutes  
and/or ions in roots. These responses should be involved in the maintenance of  
the relative water content in the leaves of grafted pepper genotypes in the  
experiment. Despite the reduction of the leaf osmotic potential and stomatal  
conductance described in the ungrafted plants, no root conductivity recovery  
should occur in the experiment since RWC was significantly lower under  
salinity. According to this relation, a reduction in either the functionality or the

1 amount of aquaporins has been reported to occur in pepper plants under  
2 salinity [30,58].  
3

4 In this experiment, the Na<sup>+</sup> and Cl<sup>-</sup> concentrations did not provoke salt  
5 toxicity symptoms in our pepper plants, and only minor leaf chlorosis and small  
6 necrotic areas was/were observed in ungrafted plants. These results agree with  
7 the lipid peroxidation levels reported in ungrafted plants when compared with  
8 grafted plants. Lower MDA concentrations were found in A/12 plants, followed  
9 by A/14. Nevertheless, gas exchange parameters were affected after a 2-week  
10 salt exposure and extended to T2. Excessive Na<sup>+</sup> and Cl<sup>-</sup> accumulation is  
11 harmful and may disrupt the integrity of the photosynthetic apparatus [24].  
12 Reduced photosynthetic capacity can be related to higher leaf Na<sup>+</sup> or Cl<sup>-</sup>  
13 concentrations [22,28,59,60]. In our experiments, the highly significant  
14 correlation found between A<sub>N</sub> and Na<sup>+</sup> and Cl<sup>-</sup> foliar concentrations suggested  
15 that both ions can be involved in reduced photosynthesis, although the  
16 regression analyses indicated a predominant inhibition effect by Na<sup>+</sup>. This  
17 effect can be linked to the concentration level in leaves and/or a major toxic  
18 power to promote inhibition. In contrast to the reductions observed in the other  
19 plant combinations, maintenance of A<sub>N</sub> in the A/12 plants can be attributed, at  
20 least in part, to increased K<sup>+</sup> levels or to other beneficial macronutrients, such  
21 as Ca<sup>2+</sup> and Mg<sup>2+</sup>, which contribute to better regulate stomata regulation under  
22 salinity [35]. Notwithstanding, g<sub>s</sub> significantly lowered under mild salt stress in all  
23 the plant combinations and for the time exposures, which corroborates a  
24 previous finding that g<sub>s</sub> are very sensitive to salt [49,61]. In addition, the  
25 diminished instantaneous carboxylation efficiency (A<sub>N</sub>/C<sub>i</sub>) noted at T1 and T2 in  
26 the ungrafted, A/5 and A/14 plants suggests that salt stress affects  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

photosynthesis by metabolic limitations, probably in association with reduced Rubisco carboxylase activity [62]. In contrast, stomatal limitations to photosynthesis should occur in A/12 at T1 and T2 since no changes in  $A_N/C_i$  were observed under salinity [63].

There is evidence that photosynthesis regulates nitrate reduction by modulating NR activity [64,65], which agrees with the results presented herein, which indicate that salt application diminishes  $A_N$  and NR activity. The most tolerant rootstock (A/12) in  $A_N$  terms exhibited lower NR inhibition if compared with the others. A drop in NR by salt can be due to: reduced nitrate transport to leaves, mainly because of nitrate/chloride competition [66]; inactivation of  $\text{NO}_3^-$  transporters by toxic effects of salt ions [67]; the disruption of root membrane integrity [58]; diminished  $\text{NO}_3^-$  transport from roots to leaves due to a lower transpiration flow [15] and, consequently, low  $\text{NO}_3^-$  loading into the root xylem, which affects NR activity [68]. Accordingly, and in accordance with the results obtained, the more marked decrease noted in NR activity (ungrafted and A/5 plants) in leaves can be associated with higher  $\text{Cl}^-$  and  $\text{Na}^+$  accumulations and/or lower carbon fixation rates.

In conclusion, the greater salt tolerance of grafted plants, mainly the A/12 (and A/14) combinations, can be attributed to their ability to restrict  $\text{Cl}^-$  transport to leaves and to diminished  $\text{Na}^+$  loading in roots, thus favouring  $\text{K}^+$  ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ ) uptake and allowing a smaller osmotic potential with a lower energy cost. These traits led to a minor inhibitory effect on photosynthesis and NR activity, which favourably affected yield [5] when compared with the A/5 and ungrafted plants. Knowledge of the physiological and biochemical processes that promote salt stress tolerance can improve our understanding of not only the mechanisms

involved in the scion and rootstock interaction, but also of the selection of robust rootstocks to be used under field salinity conditions.

## Acknowledgements

This work has been financed by INIA (Spain) through Project RTA2010-00038-C01 and the European Regional Development Fund (ERDF). C.P. is beneficiary of a doctoral fellowship (FPI-INIA).

## References

- [1] G. Colla, Y. Roupshael, C. Leonardi, Z. Bie, Role of grafting in vegetable crops grown under saline conditions, *Sci. Hortic.* 127 (2010) 147–155.
- [2] Z. Plaut, M. Edelstein, M. Ben-Hur, Overcoming salinity barriers to crop production using traditional methods, *Crit. Rev. Plant Sci.* 32 (2013) 250–291.
- [3] A. Kurunc, A. Unlukara, B. Cemek, Salinity and drought affect yield response of bell pepper similarly, *Acta Agric. Scand. Sect. B — Soil Plant Sci.* 61 (2011) 514–522.
- [4] H. Aktas, K. Abak, I. Cakmak, Genotypic variation in the response of pepper to salinity, *Sci. Hortic.* 110 (2006) 260–266.
- [5] C. Penella, S.G. Nebauer, S. López-galarza, A. Sanbautista, E. Gorbe, Evaluation for salt stress tolerance of pepper genotypes to be used as rootstocks, *J. Food, Agric. Environ.* 11 (2013) 1101–1107.
- [6] M.T. Estañ, M.M. Martinez-Rodriguez, F. Perez-Alfocea, T.J. Flowers, M.C. Bolarin, Grafting raises the salt tolerance of tomato through limiting the transport of sodium and chloride to the shoot., *J. Exp. Bot.* 56 (2005) 703–712.
- [7] M. Edelstein, Z. Plaut, M. Ben-Hur, Sodium and chloride exclusion and retention by non-grafted and grafted melon and Cucurbita plants, *J. Exp. Bot.* 62 (2011) 177–184.
- [8] Y. Huang, Z. Bie, P. Liu, M. Niu, A. Zhen, Z. Liu, et al., Reciprocal grafting between cucumber and pumpkin demonstrates the roles of the rootstock

1 in the determination of cucumber salt tolerance and sodium accumulation,  
2 Sci. Hortic. 149 (2013) 47–54.

- 3  
4 [9] F. Orsini, R. Sanoubar, G.B. Oztekin, N. Kappel, M. Tepecik, C.  
5 Quacquarelli, et al., Improved stomatal regulation and ion partitioning  
6 boosts salt tolerance in grafted melon, *Funct. Plant Biol.* 40 (2013) 628–  
7 636.  
8  
9 [10] M.M. Martinez-Rodriguez, M.T. Estañ, E. Moyano, J.O. Garcia-Abellan,  
10 F.B. Flores, J.F. Campos, et al., The effectiveness of grafting to improve  
11 salt tolerance in tomato when an “excluder” genotype is used as scion,  
12 *Environ. Exp. Bot.* 63 (2008) 392–401.  
13  
14 [11] J. Zhu, Z. Bie, Y. Huang, X. Han, Effect of grafting on the growth and ion  
15 concentrations of cucumber seedlings under NaCl stress, *Soil Sci. Plant*  
16 *Nutr.* 54 (2008) 895–902.  
17  
18 [12] S.L. Ferreira-Silva, E.N. Silva, F.E.L. Carvalho, C.S. de Lima, F.A.L.  
19 Alves, J.A.G. Silveira, Physiological alterations modulated by rootstock  
20 and scion combination in cashew under salinity, *Sci. Hortic.* 127 (2010)  
21 39–45.  
22  
23 [13] F. Giuffrida, C. Cassaniti, C. Leonardi, The influence of rootstock on  
24 growth and ion concentrations in pepper (*Capsicum annuum* L.) under  
25 saline conditions, *J. Hortic. Sci. Biotechnol.* 88 (2013) 110–116.  
26  
27 [14] C. Sonneveld, N. Straver, R. Donnan, Nutrient solutions for vegetables  
28 and flowers grown in water or substrates, Naaldwijk, The Netherlands:  
29 Proefstation voor tuinbouw onder glas te Naaldwijk. 10° ed, 1994.  
30  
31 [15] C. Penella, S.G. Nebauer, A.S. Bautista, S. López-Galarza, Á. Calatayud,  
32 Rootstock alleviates PEG-induced water stress in grafted pepper  
33 seedlings: Physiological responses, *J. Plant Physiol.* 171 (2014) 842–851.  
34  
35 [16] L.S. Bates, R.P. Waldren, I.D. Teare, Rapid determination of free proline  
36 for water-stress studies, *Plant Soil.* 39 (1973) 205–207.  
37  
38 [17] R.H. Hageman, D.P. Hucklesby, Nitrate reductase from higher plants,  
39 *Methods Enzymol.* 23 (1971) 491–503.  
40  
41 [18] E.G. Jaworski, Nitrate reductase assay in intact plant tissues., *Biochem.*  
42 *Biophys. Res. Commun.* 43 (1971) 1274–1279.  
43  
44 [19] R.L. Heath, L. Packer, Photoperoxidation in isolated chloroplasts. I.  
45 Kinetics and stoichiometry of fatty acid peroxidation, *Arch. Biochem.*  
46 *Biophys.* 125 (1968) 189–198.  
47  
48 [20] R.S. Dhindsa, P. Plumb-Dhindsa, T.A. Thorpe, Leaf senescence:  
49 Correlated with increased levels of membrane permeability and lipid  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

peroxidation, and decreased levels of superoxide dismutase and catalase, J. Exp. Bot. 32 (1981) 93–101.

- [21] R. Munns, M. Tester, Mechanisms of salinity tolerance, Annu. Rev. Plant Biol. 59 (2008) 651–681.
- [22] N. Fernández-García, V. Martínez, M. Carvajal, Effect of salinity on growth, mineral composition, and water relations of grafted tomato plants, J. Plant Nutr. Soil Sci. 167 (2004) 616–622.
- [23] L. Romero, A. Belakbir, L. Ragala, M.J. Ruiz, Response of plant yield and leaf pigments to saline conditions: effectiveness of different rootstocks in melon plant (*Cucumis melo* L.), Soil Sci. Plant Nutr. 43 (1997) 855–862.
- [24] Y. Roupahel, M. Cardarelli, E. Rea, G. Colla, Improving melon and cucumber photosynthetic activity, mineral composition, and growth performance under salinity stress by grafting onto Cucurbita hybrid rootstocks, Photosynthetica. 50 (2012) 180–188.
- [25] M. Edelstein, M. Ben-Hur, R. Cohen, Y. Burger, I. Ravina, Boron and salinity effects on grafted and non-grafted melon plants, Plant Soil. 269 (2005) 273–284.
- [26] G. Chen, X. Fu, S. Herman Lips, M. Sagi, Control of plant growth resides in the shoot, and not in the root, in reciprocal grafts of flacca and wild-type tomato (*Lysopersicon esculentum*), in the presence and absence of salinity stress, Plant Soil. 256 (2003) 205–215.
- [27] D. Savvas, A. Savva, G. Ntatsi, A. Ropokis, I. Karapanos, A. Krumbein, et al., Effects of three commercial rootstocks on mineral nutrition, fruit yield, and quality of salinized tomato, J. Plant Nutr. Soil Sci. 174 (2011) 154–162.
- [28] G. Colla, Y. Roupahel, Effect of salinity on yield, fruit quality, leaf gas exchange, and mineral composition of grafted watermelon plants, HortScience. 41 (2006) 622–627.
- [29] Y. Huang, Z. Bie, S. He, B. Hua, A. Zhen, Z. Liu, Improving cucumber tolerance to major nutrients induced salinity by grafting onto Cucurbita ficifolia, Environ. Exp. Bot. 69 (2010) 32–38.
- [30] J.M. Navarro, C. Garrido, V. Martínez, M. Carvajal, Water relations and xylem transport of nutrients in pepper plants grown under two different salts stress regimes, Plant Growth Regul. 41 (2003) 237–245.
- [31] K. Chartzoulakis, G. Klapaki, Response of two greenhouse pepper hybrids to NaCl salinity during different growth stages, Sci. Hortic. 86 (2000) 247–260.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [32] A. Altman, K. Mendel, Characteristics of the uptake mechanism of chloride ions in excised roots of a woody plant (Citrus), *Physiol. Plant.* 29 (1973) 157–162.
- [33] G. Colla, C.M.C. Suarez, M. Cardarelli, Y. Roupshael, Improving Nitrogen Use Efficiency in Melon by Grafting, *Hortscience.* 45 (2010) 559–565.
- [34] K.M. Volkmar, Y. Hu, S. H., Physiological responses of plants to salinity: a review, *Can. J. Plant Sci.* 78 (1997) 19–27.
- [35] Y. Hu, U. Schmidhalter, Drought and salinity: A comparison of their effects on mineral nutrition of plants, *J. Plant Nutr. Soil Sci.* 168 (2005) 541–549.
- [36] M.A. Hajibagheri, A.R. Yeo, T.J. Flowers, J.C. Collins, Salinity resistance in *Zea mays*: fluxes of potassium, sodium and chloride, cytoplasmic concentrations and microsomal membrane lipids, *Plant, Cell Environ.* 12 (1989) 753–757.
- [37] A. Albacete, C. Martínez-Andújar, M.E. Ghanem, M. Acosta, J. Sánchez-Bravo, M.J. Asins, et al., Rootstock-mediated changes in xylem ionic and hormonal status are correlated with delayed leaf senescence, and increased leaf area and crop productivity in salinized tomato., *Plant. Cell Environ.* 32 (2009) 928–38.
- [38] J. Gorham, Salt tolerance in the Triticeae: K/Na discrimination in some perennial wheatgrasses and their amphiploids with wheat, *J. Exp. Bot.* 45 (1994) 441–447.
- [39] A. Blum, R. Munns, J.B. Passioura, N.C. Turner, R.E. Sharp, J.S. Boyer, et al., Genetically engineered plants resistant to soil drying and salt stress: How to interpret osmotic relations?, *Plant Physiol.* 110 (1996) 1051–1053.
- [40] R. Munns, C. Brady, E. Barlow, Solute accumulation in the apex and leaves of wheat during water stress, *Aust. J. Plant Physiol.* 6 (1979) 379–389.
- [41] J. Morgan, Osmotic components and properties associated with genotypic differences in osmoregulation in wheat, *Aust. J. Plant Physiol.* 19 (1992) 67–76.
- [42] S.A. Nio, G.R. Cawthray, L.J. Wade, T.D. Colmer, Pattern of solutes accumulated during leaf osmotic adjustment as related to duration of water deficit for wheat at the reproductive stage., *Plant Physiol. Biochem.* 49 (2011) 1126–1137.
- [43] A. Patakas, N. Nikolaou, E. Zioziou, K. Radoglou, B. Noitsakis, The role of organic solute and ion accumulation in osmotic adjustment in drought-stressed grapevines, *Plant Sci.* 163 (2002) 361–367.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [44] M.C. Martínez-Ballesta, V. Martínez, M. Carvajal, Aquaporin functionality in relation to H<sup>+</sup>-ATPase activity in root cells of *Capsicum annuum* grown under salinity, *Physiol. Plant.* 117 (2003) 413–420.
- [45] S. Radić, P. Peharec Štefanić, H. Lepeduš, V. Roje, B. Pevaljek-Kozlina, Salt tolerance of *Centaurea ragusina* L. is associated with efficient osmotic adjustment and increased antioxidative capacity, *Environ. Exp. Bot.* 87 (2013) 39–48.
- [46] J.A.G. Silveira, S.A.M. Araújo, J.P.M.S. Lima, R.A. Viégas, Roots and leaves display contrasting osmotic adjustment mechanisms in response to NaCl-salinity in *Atriplex nummularia*, *Environ. Exp. Bot.* 66 (2009) 1–8.
- [47] A. Ueda, M. Kanechi, Y. Uno, N. Inagaki, Photosynthetic limitations of a halophyte sea aster (*Aster tripolium* L) under water stress and NaCl stress, *J. Plant Res.* 116 (2003) 65–70.
- [48] R. Munns, Comparative physiology of salt and water stress, *Plant Cell Environ.* 25 (2002) 239–250.
- [49] Y. He, Z. Zhu, J. Yang, X. Ni, B. Zhu, Grafting increases the salt tolerance of tomato by improvement of photosynthesis and enhancement of antioxidant enzymes activity, *Environ. Exp. Bot.* 66 (2009) 270–278.
- [50] J.M. Ruiz, J.J. Ríos, M.A. Rosales, R.M. Rivero, L. Romero, Grafting between tobacco plants to enhance salinity tolerance, *J. Plant Physiol.* 163 (2006) 1229–1237.
- [51] T.D. Colmer, R. Munns, T.J. Flowers, Improving salt tolerance of wheat and barley: future prospects, *Aust. J. Exp. Agric.* 45 (2005) 1425–1443.
- [52] Z. Chen, T.A. Cuin, M. Zhou, A. Twomey, B.P. Naidu, S. Shabala, Compatible solute accumulation and stress-mitigating effects in barley genotypes contrasting in their salt tolerance, *J. Exp. Bot.* 58 (2007) 4245–4255.
- [53] S. Lutts, G. Guerrier, Peroxidase activities of two rice cultivars differing in salinity tolerance as affected by proline and NaCl, *Biol. Plant.* 37 (1995) 577–586.
- [54] M.W. Hester, I.A. Mendelsohn, K.L. McKee, Species and population variation to salinity stress in *Panicum hemitomom*, *Spartina patens*, and *Spartina alterniflora*: Morphological and physiological constraints, *Environ. Exp. Bot.* 46 (2001) 277–297.
- [55] J. Raven, Regulation of pH and generation of osmolarity in vascular plants: a cost-benefit analysis in relation to efficiency of use of energy, nitrogen and water, *New Phytol.* 101 (1985) 25–77.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [56] A. Poljakoff-Mayber, Morphological and anatomical changes in plants as a response to salinity stress, A Poljakoff-Mayber and J Gale (eds). Plants in saline environments. Ecological Series 15, Springer-Verlag, Berlin, Germany, 1975.
- [57] M. Calvo-Polanco, B. Sánchez-Romera, R. Aroca, Mild salt stress conditions induce different responses in root hydraulic conductivity of *Phaseolus vulgaris* over-time, PLoS One. 9 (2014). 3e90631.
- [58] M. Carvajal, V. Martínez, C.F. Alcaraz, Physiological function of water channels as affected by salinity in roots of paprika pepper, Physiol. Plant. 105 (1999) 95–101.
- [59] R.R, Walker, D.H. Blackmore, Q. Sung, Carbon dioxide assimilation and foliar ion concentration in leaves of lemon (*Citrus limon* L.) trees irrigated with NaCl or Na<sub>2</sub>SO<sub>4</sub>. Aust. J. Plant Physiol. 20 (1993) 173-175.
- [60] M.C. Martínez-Ballesta, V. Martínez, M. Carvajal, Osmotic adjustment, water relations and gas exchange in pepper plants grown under NaCl or KCl, Environ. Exp. Bot. 52 (2004) 161–174.
- [61] Q. Jiang, D. Roche, T.A. Monaco, D. Hole, Stomatal conductance is a key parameter to assess limitations to photosynthesis and growth potential in barley genotypes, Plant Biol. 8 (2006) 515–521.
- [62] E.N. da Silva, R.V. Ribeiro, S.L. Ferreira-Silva, R.A. Viégas RA, J.A.G. Silveira, Salt stress induced damages on the photosynthesis of physic nut young plants, Sci. Agric. 68 (2011) 62-68.
- [63] J. Flexas, J. Bota, F. Loreto, G. Cornic, T.D. Sharkey, Diffusive and metabolic limitations to photosynthesis under drought and salinity in C) plants, Plant Biol. 6 (2004) 269–79.
- [64] W.M. Kaiser, E. Brendle-Behnisch, Rapid Modulation of Spinach Leaf Nitrate Reductase Activity by Photosynthesis: I. Modulation *in vivo* by CO<sub>2</sub> Availability, Plant Physiol. 96 (1991) 363–367.
- [65] S. Yousfi, M.D. Serret, A.J. Márquez, J. Voltas, J.L. Araus, Combined use of  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$  and  $\delta^{15}\text{N}$  tracks nitrogen metabolism and genotypic adaptation of durum wheat to salinity and water deficit, New Phytol. 194 (2012) 230–44.
- [66] G.K. Abd-El-Baki, F. Siefritz, H.M. Man, H. Weiner, R. Kaldenhoff, W.M. Kaiser, Nitrate reductase in *Zea mays* L. under salinity, Plant, Cell Environ. 23 (2000) 515–521.
- [67] H. Lin, S.S. Sandra, K.S. Schumaker, Salt sensitivity and the activities of H-ATPase in cotton seedlings, Crop Sci. 37 (1997) 190-197.

1 [68] M. Debouba, H. Maârroufi-Dghimi, A. Suzuki, M.H. Ghorbel, H. Gouia,  
2 Changes in growth and activity of enzymes involved in nitrate reduction  
3 and ammonium assimilation in tomato seedlings in response to NaCl  
4 stress, Ann. Bot. 99 (2007) 1143–1151.  
5  
6  
7  
8  
9

## 10 Legends of figures

11  
12  
13  
14 **Fig. 1.** Effect of NaCl addition at 0 mM (□) and 40mM (■) on relative  
15 leaf water content (RWC %) for exposures of 14 days (A) and 28 days (B) in  
16 ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto  
17 accessions 5, 12 and 14. Dates are mean values±SE for n=6. In each plant  
18 combination, different letters indicate significant differences at  $P < 0.05$  (LSD  
19 test).  
20  
21  
22  
23  
24  
25  
26  
27

28  
29  
30  
31 **Fig. 2.** Leaf osmotic potential (MPa) in ungrafted pepper plants (cultivar 'Adige')  
32 and the cultivar grafted onto accessions 5, 12, and 14 after addition of NaCl at  
33 0mM (□) and 40mM (■) for exposures of 14 days (A) and 28 days (B).  
34 Dates are mean values±SE for n=6. In each plant combination, different letters  
35 indicate significant differences at  $P < 0.05$  (LSD test).  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 **Fig.3.** Concentrations of Na<sup>+</sup> (A, B), Cl<sup>-</sup> (C, D), K<sup>+</sup> (E, F) in mg g<sup>-1</sup> DW and the  
47 Na<sup>+</sup>/K<sup>+</sup> ratio (G, H) in the leaves and roots of ungrafted pepper plants (cultivar  
48 'Adige') and the cultivar grafted onto accessions 5, 12, and 14 after addition of  
49 NaCl at 0mM and 40mM for exposures of 14 days (□, ■) and 28 days  
50 (▨, ■), respectively. Dates are mean values±SE for n=6. In each  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

**Fig. 4.** Ionic concentration of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  in the leaves (A, C) and roots (B, D) in  $\text{mg g}^{-1}$  DW of ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto accessions 5, 12, and 14 after addition of NaCl at 0mM and 40mM for exposures of 14 days (□, ■) and 28 days (▨, ▩), respectively. Dates are mean values $\pm$ SE for  $n=6$ . In each plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

**Fig. 5.** Changes in the proline concentration ( $\text{mg proline g}^{-1}\text{DW}$ ) from ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto accessions 5, 12 and 14 after addition of NaCl at 0mM (□) and 40mM (■) for exposures of 14 days (A) and 28 days (B). Dates are mean values $\pm$ SE for  $n=6$ . In each plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

**Fig. 6.** The Net  $\text{CO}_2$  assimilation rate ( $A_N$ ;  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) (A, B); leaf stomatal conductance ( $g_s$ ;  $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ ) (C, D) and instantaneous carboxylation efficiency ( $A_N/C_i$ ; E, F) in ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto accessions 5, 12 and 14 after addition of NaCl at 0mM (□) and 40mM (■) for exposures of 14 days (A, C, E) and 28 days (B, D, F). Dates are mean values $\pm$ SE for  $n=10$ . In each plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Fig. 7.** Nitrate reductase activity ( $\mu\text{mol NO}_2 \text{ g}^{-1} \text{ FW h}$ ) in the leaves of ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto accessions 5, 12 and 14 after addition of NaCl at 0mM (□) and 40mM (■) for exposures of 14 days (A) and 28 days (B). Dates are mean values $\pm$ SE for n=6. In each plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

**Fig. 8.** Leaf malondialdehyde (MDA) content ( $\text{nmol MDA g}^{-1} \text{ FW}$ ) in the leaves of ungrafted pepper plants (cultivar 'Adige') and the cultivar grafted onto accessions 5, 12 and 14 after addition of NaCl at 0mM (□) and 40mM (■) for exposures of 14 days (A) and 28 days (B). Dates are mean values $\pm$ SE for n=6. In each plant combination, different letters indicate significant differences at  $P < 0.05$  (LSD test).

**Table 1.**

Linear regression and statistical analysis between mineral ions concentration (mg g<sup>-1</sup> DW) in the leaves of the cultivar “Adige” ungrafted and grafted onto different pepper genotypes (5, 12 and 14), and proline (mg g<sup>-1</sup> DW), as related to the osmotic potential ( $\Psi_s$  s in MPa) and CO<sub>2</sub> fixation rate ( $A_N$ ,  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ).

Salt treatment time	Regression equations	$P^*$	$R^2$
T1	$\Psi_s = -0.021[\text{Na}^+] - 1.05$	0.0035	0.782
	$\Psi_s = -0.006 [\text{Cl}^-] - 0.99$	0.0003	0.898
	$\Psi_s = 0.02 [\text{K}^+] - 1.61$	0.008	0.716
	$\Psi_s = -0.22 [\text{Proline}] - 0.64$	0.0229	0.616
	$A_N = -0.641 [\text{Na}^+] + 21.35$	0.0002	0.919
	$A_N = -0.1616 [\text{Cl}^-] + 22.47$	0.0017	0.828
	$A_N = 0.642 [\text{K}^+] + 2.48$	0.1333 ns	0.701
	$A_N = -7.856 [\text{Proline}] + 37.28$	0.0548 ns	0.593
T2	$\Psi_s = -0.029 [\text{Na}^+] - 1.13$	0.0001	0.923
	$\Psi_s = -0.0065 [\text{Cl}^-] - 1.105$	0.0031	0.792
	$\Psi_s = 0.021 [\text{K}^+] - 1.87$	0.0514 ns	0.495
	$\Psi_s = -0.143 [\text{Proline}] - 1.02$	0.0332	0.986
	$A_N = -0.647 [\text{Na}^+] + 21.81$	0.0004	0.897
	$A_N = -0.144 [\text{Cl}^-] + 22.46$	0.0032	0.788
	$A_N = 0.537 [\text{K}^+] + 4.88$	0.0794 ns	0.635
	$A_N = -2.943[\text{Proline}] + 24.38$	0.0018	0.910

Determinations were made after 14 (T1) and 28 (T2) days. Fisher's least significance difference (LSD) test at  $P < 0.05$  was used.

\*For all the linear regressions, the degrees of freedom are  $n = 4$ .

Figure 1







Figure 2






Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

